





An Agile Development Cycle of an Online Memory Program for Healthy Older Adults

www.cambridge.org/cjg

Iris Yusupov^{1,2} , Susan Vander Morris¹ , Cindy Plunkett^{1,3}, Arlene Astell^{4,5,6} , Jill B. Rich^{1,2} and Angela K. Troyer^{1,4} 

Research Note / Note de recherche

Cite this article: Yusupov I, Vander Morris S, Plunkett C, Astell A, Rich JB, & Troyer AK. (2022). An Agile Development Cycle of an Online Memory Program for Healthy Older Adults. *Canadian Journal on Aging / La Revue canadienne du vieillissement*
<https://doi.org/10.1017/S0714980821000763>

Received: 27 September 2020

Accepted: 12 July 2021

Mots-clés:

vieillesse; en ligne; mémoire; intervention; personnes âgées; développement de programmes; apprentissage en ligne

Keywords:

aging; online; memory; intervention; older adult; program development; e-learning

Corresponding author:

La correspondance et les demandes de tirés-à-part doivent être adressées à : /
Correspondence and requests for offprints should be sent to: Iris Yusupov, M.A., Behavioural Science Building, Psychology Department, York University, 4700 Keele St., Toronto, ON M3J 1P3, Canada (iris@yorku.ca)

¹Baycrest Health Sciences, North York, Ontario, Canada, ²Department of Psychology, York University, Toronto, Ontario, Canada, ³PointClickCare, Mississauga, Ontario, Canada, ⁴Department of Psychiatry, University of Toronto, Toronto, Ontario, Canada, ⁵University Health Network, Toronto, Canada and ⁶University of Reading, Reading, UK

Résumé

Les interventions en ligne pour les personnes âgées doivent être adaptées à leurs besoins spécifiques afin d'augmenter leur efficacité et l'adhésion des utilisateurs. Le cycle de développement agile est un modèle dynamique permettant de solliciter et d'intégrer les commentaires des personnes âgées au cours du processus de conception. Nous avons combiné cette approche avec le Cadre des phases cliniques et translationnelles de l'université Harvard qui fournit une structure claire pour évaluer les nouveaux programmes de santé avant qu'ils ne soient proposés dans la communauté. Nous avons élaboré notre programme en ligne sur la mémoire à partir du programme sur la mémoire et le vieillissement qui se donnait en présentiel, et qui avait été validé empiriquement. L'objectif de l'étude était d'associer le cycle de développement agile avec le Cadre des phases cliniques et translationnelles pour concevoir et tester un programme pilote en ligne sur la mémoire qui est adapté aux besoins uniques des personnes âgées. L'étude no 1 a permis de tester dans une phase pilote les modules individuels du programme avec des participants présents sur place, et à intégrer leurs commentaires lors du développement du programme afin d'en optimiser la convivialité. L'étude no 2 a consisté en deux pilotes séquentiels du programme accessibles à distance dans lesquels les résultats cliniques préliminaires ont été évalués et des commentaires ont été collectés pour guider les modifications itératives. Les plans pour une validation ultérieure et les limites des études sont discutés. L'application réussie du cycle de développement agile mis en œuvre dans cette série d'études pourra être adaptée par d'autres équipes souhaitant proposer un contenu en ligne à des groupes d'utilisateurs finaux plus spécifiques.

Abstract

Online interventions for older adults should be tailored to their unique needs to increase the efficacy of and adherence to the intervention. The agile development cycle is a dynamic model to solicit and incorporate feedback from older adults during the design process. We combined this approach with the framework of Harvard University's clinical and translational phases that provide a clear structure for evaluating new health programs before they are offered in the community. We based our online memory program on the empirically validated in-person Memory and Aging Program. The aim of the present study was to combine the agile development cycle with the clinical and translational phases framework to develop and pilot an online memory program tailored to the unique needs of older adults. Study 1 involved piloting individual program modules on site and integrating participant feedback into the program's design to optimize usability. Study 2 involved two sequential pilots of the program accessed remotely to evaluate preliminary clinical outcomes and obtain feedback for iterative modifications. Plans for further validation and limitations are discussed. The successful application of the agile development cycle implemented in this series of studies can be adapted by others seeking to offer online content for targeted end users.

The majority of older adults will experience normal age-related cognitive changes (Craik & Rose, 2012), for which an increasing number of training programs have been created to optimize memory functioning. Research shows that memory interventions can improve performance on trained memory tasks (Karbach & Verhaeghen, 2014), increase knowledge about memory, and promote healthy lifestyle changes to support brain health (Vander Morris, Au, Gardner, & Troyer, 2020), and bolster self-reported memory abilities and strategy use in daily life (Hudes, Rich, Troyer, Yusupov, & Vander Morris, 2019).

Although many older adults who experience cognitive changes are proactively looking for ways to keep their brain active (Parikh, Troyer, Maione, & Murphy, 2016), there are several limitations to doing this through *in-person* intervention programs. Participation may be

© Canadian Association on Gerontology 2022.

CAMBRIDGE
UNIVERSITY PRESS

restricted for some individuals for a variety of reasons such as difficulties traveling to the site because of physical disabilities, limited access to transportation, or living in remote areas (Fitzpatrick, Powe, Cooper, Ives, & Robbins, 2004; Pike et al., 2018) or cancellation of in-person group activities, as seen during the COVID-19 pandemic. In response to the pandemic, the already increasing rate of technology use among older adults has surged, with more than 88 per cent of Canadians over the age of 65 using the Internet daily (AGE-WELL, 2020). More specifically, although this is an emerging field, research shows that older adults do benefit from online cognitive programs (for a review, see Kueider, Parisi, Gross, & Rebok, 2012). Therefore, creating an online memory intervention may increase accessibility to a broader audience and offer greater convenience, privacy, and flexibility in scheduling for participants.

In considering an online program, it is important to acknowledge that online interventions often have higher rates of attrition than in-person interventions. For example, a systematic review of 83 online health interventions found an adherence rate of approximately 50 per cent (Kelders, Kok, Ossebaard, & van Gemert-Pijnen, 2012). Personalization of online interventions is essential for promoting participant engagement and satisfaction, thus leading to greater adherence and efficacy of the given program. One avenue to achieve personalization is to develop an online program for a target group and to effectively understand the target group's needs during the design process (Ludden, van Rompay, Kelders, & van Gemert-Pijnen, 2015). In a recent review of online memory programs for older adults, Pike et al. (2018) offer practice recommendations for their development including considering age-associated sensory and cognitive changes in the program's design, while ensuring that the program is easy to learn and intuitive. However, no specific recommendations are provided to inform the development process and its piloting in order to gauge usability.

Agile Development Cycle

In this article, we describe the successful application of an agile development cycle to the creation of an online memory intervention program for older adults. An agile development cycle is an iterative process adapted from the technology sector that incorporates end users' feedback during each phase of program development and piloting (Davis, 2013). This approach is in contrast with the traditional Waterfall Models (e.g., the ADDIE model which involves the Analysis, Design, Development, Implementation, and Evaluation stages) for instructional design and development, which are essentially linear, inflexible approaches that require the completion of each phase prior to moving on to the next (Stoica, Ghilic-Micu, Mircea, & Uscatu, 2016). The assumption underlying Waterfall Models is that developers possess all the necessary information for the designing process (Doolittle, 2020). An agile development cycle, in contrast, is better suited for developing online learning programs for specific populations (i.e., older adults) because it allows for tailoring the program to their unique needs, which may not be fully known or understood at the onset. It also allows for flexibility in making modifications to preceding phases of design (Davis, 2013; see Figure 1).

Translational Phases

Considering that the agile development cycle is adapted from the technology sector, we combined this process with the framework from the Harvard Clinical and Translational Science Center's five




Figure 1. Illustration of the traditional Waterfall Model in contrast to the agile development cycle, which has the fluid capability to return to preceding phases of testing and development.

translational (T) phases that are necessary to execute before a health intervention can become available to the general public (Harvard Catalyst, 2021). In the next section, we describe the foundational work of the first two phases, which involve seeking evidence from the extant literature (T0) and proof of concept for the online intervention (T1). The current research focuses on the subsequent phases, which involves testing the intervention in a controlled environment (T2) and testing the intervention in the intended environment of the final product (i.e., remotely accessed) while assessing preliminary program outcomes (T3). The final phase evaluating benefits under a more rigorous research design (T4) is discussed as a future step in the discussion. See Figure 2 for an infographic of the phases, based on material from Sung et al. (2003), Szilagyi (2009), and Westfall, Mold, and Fagnan (2007).

Foundational Work

T0: Basic Research

The first translational phase, T0, involves understanding the biopsychosocial mechanisms of an underlying health problem and seeking opportunities for its treatment (Harvard Catalyst, 2021). As reviewed, older adults can benefit from carefully crafted memory programs grounded in research on effective memory strategies and modifiable lifestyle factors. Further, Pike et al. (2018) recommend considering existing and effective in-person memory programs when developing an online program. Based on this literature, we chose to develop an online program modelling an empirically validated in-person memory intervention called The Memory & Aging Program[®], which has been offered for more than 20 years at Baycrest, a global leader in geriatric research and care in Toronto (Troyer, 2001). This program aligns with practice recommendations by Pike et al., as it offers its participants psychoeducation on aging and memory, a “tool kit” of memory strategies, and group discussions (Troyer & Vander Morris, 2012).

Benefits of the Memory and Aging Program include increased memory knowledge and strategy use, increased satisfaction and confidence with one's everyday memory functioning (Troyer, 2001), positive change in healthy lifestyle domains (Vander Morris et al., 2020), decrease in intentions to seek unnecessary medical attention for memory concerns (Wiegand, Troyer, Gojmerac, & Murphy, 2013), and feelings of acceptance and




Figure 2. Overview of the Harvard Clinical and Translational Science Center's five translational phases; adapted framework for the testing of a health intervention prior to its release to the general public.

reduced anxiety about normal age-related memory changes (Vandermorris et al., 2017).

T1: Translation to Humans

The second translational phase, T1, involves proof of concept and understanding the feasibility of translating the original in-person intervention into an online format (Harvard Catalyst, 2021). A multidisciplinary team was created that included research and clinical neuropsychologists (A.K.T., S.V.) and instructional design and e-learning development experts (including C.P.). Patient advisors (e.g., graduates of the in-person program) also provided input.

The multidisciplinary team used program materials for the in-person program (Troyer & Vandermorris, 2012; 2017) and followed the next steps for creating a new e-learning program, including:

1. *Action Mapping*, which began with assigning roles within the e-learning team, discussing delivery method of material, and creating a timeline for task deadlines. This step focused on solving performance problems and setting measurable learning outcomes for the program.
2. *Storyboarding*, which involved creating a storyboard template, with all content reviewed by the team. Next, design elements such as themes, colour scheme, narration, and interactions were discussed and confirmed. A delivery date was set.
3. *Design and Development*, which involved taking all information determined in the *Action Mapping* and *Storyboarding* stages and

applying instructional design and development for online learning best practices to produce the e-learning program.

The initial version of the program was developed (see Table 1 for a brief description of each module) and based on themes that emerged from brainstorming sessions and patient advisor feedback, certain elements were included. For example, as privacy of personal information is a significant concern for older adults (Chang, McAllister, & McCaslin, 2015), participants were provided with the option of using a non-identifying username as opposed to their real names. They were also informed that information would remain private within the program and accessible only to other registered participants.

Current Study

Based on the foundational T0 and T1 work described, we used an agile development cycle to implement and evaluate translational phases T2 and T3, which are reported separately in the following two studies. Study 1 involved piloting individual program modules on site and integrating participant feedback into the program's design to optimize usability, with the primary aim of executing the T2 translational phase. Study 2 involved two sequential pilots of the program accessed remotely to evaluate preliminary clinical outcomes that have been demonstrated in the in-person Memory and Aging Program. The primary aim of Study 2 was to successfully execute the T3 translational phase and to continue making

Table 1. Description of individual modules in the online Memory and Aging Program

	Module Description
Getting Started	Navigating the learning management system and operating program functions
Module 1: Introduction	Overview of content and introduction to different components of the program (e.g., discussion boards, surveys, and polls)
Module 2: What is Memory?	Explanation of what memory is, what brain regions are involved in memory, types of memory processes, normal and abnormal memory changes
Module 3: Factors Affecting Memory	A discussion of health and lifestyle factors that affect memory
Module 4: Stress and Relaxation	A discussion of the effect of stress on memory and introduction of relaxation techniques
Module 5: Memory Strategies Overview and Practice Retrieval	An explanation of the rationale, procedures, and evidence supporting 5 key memory strategies, as well as in-depth practice of the spaced-retrieval memory strategy
Module 6: Memory Strategies Associations and Records	In-depth practice of the semantic association memory strategy and a discussion of the use of records as a memory aid
Module 7: Application of Memory Strategies and Goal Setting	Review of memory strategies and training on how to set effective goals
Module 8: Wrap-up and Feedback	Engaging participants in a review game, goal setting, sharing final thoughts, and providing feedback

modifications using participant feedback as per the agile development cycle.

Study 1: Translation to Patients

Methods

In this T2 phase, the intervention program was tested in a highly controlled environment (i.e., on site in a computer laboratory; Harvard Catalyst, 2021).

Participants

A total of 25 participants were involved in this phase, including local community-dwelling older adults recruited through an e-mail advertisement from a pool of hospital volunteers unfamiliar with the Memory and Aging Program ($n = 21$) and through direct requests to individuals who previously completed the in-person Memory and Aging Program ($n = 4$).

Procedure and Measures

Participants provided written informed consent for audio recording of spoken feedback and photography of their engagement with the technology. The essence of the agile development cycle involves breaking down the larger intervention into smaller cycles

(known as Sprints) and offering incremental delivery of the intervention components (Flewelling, 2018). Therefore, of these 25 participants, 14 piloted Modules 2, 3, and 5; 7 piloted Modules 4, 6, and 7; and 4 (the graduates of the in-person program) piloted Module 8. Feedback from the first pilot (Modules 2, 3, and 5) was discussed within the multidisciplinary team, and modifications were made to the proceeding modules prior to the subsequent piloting cycles. This process was repeated after each piloting session.

Clinical, research, and e-learning staff observed the participants as they engaged in the modules, paying special attention to their reactions and noting any areas of confusion and particular enjoyability. Participants filled out feedback questionnaires (The Scavenger Hunt measure from the TUNSGTEN tools: <http://tungsten-training.com>) for each module that assessed usability of new technologies in interactive settings by indicating *yes* or *no* to the following statements: (a) This is easy to use, (b) This is something I would use, and (c) This is something I enjoy (Astell, Dove, Morland, & Donovan, 2020). The questionnaire also encouraged participants to add general comments or suggestions for improvement, to enable technologies to be modified to better meet the user's needs. These were used as the guiding structure for the focus groups that followed.

Results

Piloting of Modules 2, 3, and 5 (Understanding Memory, Modifiable Lifestyle Factors, and Memory Strategies)

Piloting of these modules revealed that all 14 participants found at least one of the three modules easy to use, enjoyable, and something that they would use outside of the laboratory setting. Seven of 14 participants reported such feelings about all three of the piloted modules. Module 5, which involved an interactive component, was reportedly enjoyed by all but one participant. For example, one participant reported, "This module works well. Interactive and fun. Nice customization. Well done."

Feedback was reviewed, and several rounds of modifications were made. In general, there appeared to be difficulties with navigating the modules, as some procedures that may seem intuitive to an avid technology user were not obvious to the participants, including how to adjust the sound, pause the videos, or select items.

Piloting of Modules 4, 6, and 7 (Stress and Relaxation, Practicing Memory Strategies, and Strategies Overview)

Information gained from the first series of piloting previously detailed was incorporated into the design of Modules 4, 6, and 7 prior to in-person piloting. Responses revealed that all seven participants from this pilot session found at least one of the three modules easy to use, enjoyable, and something that they would use outside of the laboratory setting, and four participants found that all three piloted modules met these criteria.

Piloting of Module 8 (Summary and Wrap-up)

Module 8 consists of a review game, final thoughts, and creating a plan for memory or health improvement (i.e., setting goals). These components were piloted by graduates of the in-person Memory and Aging Program, as they possessed the background information requisite to participate in the review game and provide meaningful feedback on the way that the course content was summarized. With the exception of one participant who noted "maybe" to the questions about enjoyment and using the review game in a real-life setting, all four participants reported that each of the three

components of Module 8 was easy to use, enjoyable, and something that they would use outside of the laboratory setting. Specific feedback included, “Good practice run for refresher of what I learned in the program.”

Qualitative feedback

Qualitative feedback (i.e., from the open-ended questionnaire responses and focus group discussion) from all 25 participants identified a number of technological glitches and settings that needed adjustment. Subsequent modifications based on this feedback involved removing background music and excessive sound effects, removing the requirement for double-clicking, increasing font size, adjusting speed and volume of speech, and inserting additional prompts and instructions on how to proceed (e.g., a pop-up indicating “click ‘next’”). See Table 2 for more detailed participant feedback.

Study 2: Translation to Practice

Methods

Study 2 involved testing the intervention in its entirety in the intended environment for the final product (i.e., accessed remotely by participants within their community) in two separate pilot studies. In addition, we tested preliminary clinical outcomes by incorporating targeted measures that have been shown to improve following participation of the in-person program characteristic of the T3 Translation to Practice phase (Harvard Catalyst, 2021). The results of each pilot are reported separately as Sub-study 2a and Sub-study 2b (with the exception of the results from the Memory Toolbox questionnaire and program-specific goal attainment, which were pooled across both pilots and reported under Sub-study 2b).

Participants

Forty community-dwelling older adults in Canada were recruited through e-mail advertisements to participate remotely (22 participated in the first pilot and 18 participated in the second pilot). In light of a recent study that found that less frequent computer usage predicted attrition in the initial phases of online studies (Rübsamen, Akmatov, Castell, Karch, & Mikolajczyk, 2017) and in the hope of mitigating participant distress during the early stages of piloting, participants were required to self-report using a computer *at least once a day* and to feel *comfortable* or *very comfortable* using a computer.

Procedure

Participants completed online questionnaires and a telephone interview prior to the intervention (detailed subsequently). Participants were then e-mailed a link to register for the online Memory and Aging Program. Once registered, they were able to complete the pre-intervention questionnaires and watch the introduction videos. Each module was released on a weekly basis as long as the participant had completed all tasks in the previous module. The e-learning team monitored the completion of individual items for each participant. Once it was apparent that participants had completed the program (i.e., all of the modules described in Table 1), they filled out post-intervention questionnaires online and completed an interview over the telephone.

Table 2. Overview of Study 1 feedback and areas of modification

Area of Modification	Participant Feedback
Removed background music and excessive sound effects	“Don’t like music, too loud. Hard to hear dialogue.” ^a
Adjusted speed and volume of speech	“The commentary is too fast.” “Speed made this a little harder.”
Increased font size and adjusted visuals	“Progress bar should be a colour - not white on white.” <i>Font size is too small in certain slides.</i> ^b
Removed technical glitches and spelling errors	“Slight hang up before poll results opened.” “Typo in slides - ‘any’.”
Removed all double-clicking	<i>It is difficult to double-click fast enough to initiate the command.</i>
Located areas to include prompts on how to proceed or additional instructions	“No suggestion to ‘click next’.” “You should be prompted to click on ‘Next’.” “At the end no indication on how to proceed.” “Clear indication of end of a section would be helpful.” “‘Type your answers in space provided’ you must click in text box to start.” (Including instructions to “click here in order to type.”)
Formatted and modified activities	“Not enough space to complete answers.” “Would be a good idea to see the results of all questions... and maybe try to do all exercises again.” <i>I would like the opportunity to replay the review game to improve my score.</i> <i>The tone you hear when you make a mistake in the game is too discouraging.</i>
Informed the need for supplemental materials	“The directions were easy to follow, but I would need either a written manual or be able to access the directions in the days to follow.” “I don’t have a printer at home, so I would not be able to print out the materials.”

Note.^aFeedback in quotations is extracted from the written feedback questions.

^bInformation written in italics represents gist themes reported from the audio-recorded focus group discussions.

Measures

In order to gauge the program’s benefits, participants completed the following memory-related measures before and after completion of the intervention.

The Memory Knowledge Quiz. During the telephone interview, participants were asked 12 open-ended questions related to memory knowledge and brain health. Each answer was scored from 1 to 6 points, depending on the demand of the question. The Memory Knowledge Quiz was adapted from previous evaluations of the in-person version of the Memory and Aging Program (Troyer, 2001).

The Memory Toolbox. This online questionnaire provided participants with six familiar scenarios (e.g., learning the name of a new acquaintance), for which they were required to list memory strategies that would be useful for each situation (Troyer, 2001).

The number and quality of memory strategies were analyzed before and after the program.

Lifestyle behavior change. As part of the online questionnaires, a subset of participants ($n = 10$) were asked about their lifestyle behaviors with the following question: “Have you made any lifestyle changes in the past month that may improve your health and possibly memory (e.g., lower stress levels, use of relaxation techniques, improved diet or exercise, engagement in cognitively or socially engaging activities)?”

Program-specific goal attainment and general feedback. Similar to the in-person program, participants were asked to choose three program-specific goals that best aligned with their intentions for the program prior to its start (e.g., to understand how lifestyle factors affect memory, to feel more confident about their memory). During the post-intervention telephone interview, participants were reminded of their three goals and were asked to rate them on a five-point satisfaction scale. General feedback related to the program and suggestions for improvement were collected during the telephone interview and in an online questionnaire.

Results

Sub-study 2a

Of the 22 recruited participants, 11 completed all modules and the post-intervention telephone interview; 10 of these 11 participants also completed the online post-intervention questionnaires. Several areas of technical difficulty including registration, password creation, and browser compatibility were reported through e-mail communications to the e-learning team.

Memory knowledge and strategy use. For the 11 participants who completed the Memory Knowledge Quiz, scores showed a large, significant improvement from before to after program completion $t(10) = -5.85, p < 0.001, d = 1.76$. Individual scores are shown in Figure 3.

Ten participants completed the Memory Toolbox questionnaire. In the post-intervention responses, one participant applied three new memory strategies, three applied two new memory

strategies, and six applied one new memory strategy. Therefore, all 10 participants learned at least one new memory strategy that they could apply in real-life scenarios. These data were combined with data from Sub-study 2b for statistical analysis and are reported in the Sub-study 2b results section.

Lifestyle behavior change. Participants were also asked about changes in their lifestyle behaviors (e.g., use of relaxation techniques, improved diet or exercise, engagement in cognitively or socially engaging activities). Five of the 10 participants reported that they did not make any lifestyle changes in the month prior to the commencement of the online program, but that they had made a lifestyle change following program completion. Two participants indicated they had made a change within the month prior to as well as in the month following completion of the program. The remaining three participants reported before and after the program that they had not made a lifestyle change in the previous month. Based on 20 per cent of individuals making a lifestyle change at baseline, a Chi square analysis revealed that a significant proportion of participants (70%) reported a lifestyle change after completing the program ($\chi^2 = 25, df = 1, p < 0.001$).

General feedback

Through post-intervention telephone interviews and online questionnaires, participants provided general feedback and favourable ratings for their program goals (reported under Sub-study 2b). Overall, they expressed enjoyment of the various presentation formats (i.e., videos, animations, games), which kept them engaged throughout the modules. Feedback was generally positive regarding the interactive nature of the online program, such as the use of real-world examples and cartoon animations that depict common and relatable scenarios. Some participants also mentioned the usefulness of having a transcript of each slide in order to follow along with the audio component. Finally, participants expressed feelings of normalization about their memory ability.

Participants who experienced technical difficulties were able to e-mail the project coordinator and receive support from the e-learning team; thus, all areas of difficulty were systematically




Figure 3. Pre- and post-intervention individual participant scores on the Memory Knowledge Quiz during Sub-study 2a in the T3 Translation to Practice phase. Note. $n = 11$, maximum possible score of 20.

logged. Subsequently, a “Getting Started” module was added to include an introduction to various program features (such as accessing the transcript, adjusting volume, and posting in the discussion forum). In addition, a Frequently Asked Questions (FAQ) document was compiled, containing all of the areas of concern or difficulty that had been recorded.

For the homework assignments, individual exercise sheets were available for participants to print and utilize. However, some participants reported that they did not have access to a printer. This is in line with the feedback from the T2 Translation to Patients phase involving the in-person Memory and Aging Program graduates who spoke about the utility of having a participant workbook. Another subject that participants mentioned was the need for fostering greater interaction. Although there were discussion boards, there was little back and forth conversation amongst participants. In order to increase participant interaction, weekly “coffee breaks” were held as a live chat room for the participants and the facilitator to interact with each other.

Sub-study 2b

Because of the flexible and iterative nature of the agile development cycle, we returned to making modifications to individual modules by integrating the feedback from Sub-study 2a prior to launching Sub-study 2b. The goal of Sub-study 2b was to ensure that the technical glitches were resolved, obtain feedback regarding the addition of the participant workbook (mailed to the participants’ homes) and the “coffee break” chat rooms, and continue assessing program outcomes. Of the 18 participants, 9 completed the online Memory and Aging Program as well as the post-intervention telephone interview to obtain feedback. Seven of these participants also completed the online version of the Memory Toolbox questionnaire, which was administered to continue to monitor the

benefits of the program content itself and participant engagement with material, and 11 provided ratings for their program goals.

Memory Toolbox. Results of the pre- and post-intervention Memory Toolbox questionnaire indicated that all but one participant acquired and applied a new memory strategy; three out of seven added three strategies, two out of seven added two strategies, and one participant added one new strategy (see Figure 4). Across Studies 2a and 2b, participants were able to provide a significantly greater number of strategies for the given scenarios after program completion, $t(16) = -3.21, p < 0.001, d = 0.78$. An examination of the types of responses revealed a particular acquisition of internal strategies, such as paying close attention and forming implementation intentions (e.g., saying “I am turning off the stove” when doing so). In contrast, external strategies, such as keeping a record book or agenda, were most frequently listed as useful strategies by participants prior to participation in the online program.

Program-specific goal attainment. Across Sub-studies 2a and 2b, a total of 20 participants provided a rating for their three personal goals. Overall, all participants ($n = 20$) were at least *somewhat satisfied* with at least one of their chosen goals; 16 participants were at least *mostly satisfied* with at least one of their goals, and 7 participants were *completely satisfied* with at least one of their goals (see Figure 5). The large majority of goals (54 out of 60) were rated positively, indicating that participants were getting what they wanted out of the program.

General feedback

Through the post-intervention telephone interview and online questionnaires, all but one of the participants reported feeling *satisfied* with the participant workbook. This participant mentioned that the workbook was an unnecessary addition, as links to downloadable homework logs were available. Several participants




Figure 4. Pre- and post-intervention individual participant scores on the Memory Toolbox questionnaire during Sub-studies 2a and 2b in the T3 Translation to Practice phase. Note. $n = 17$




Figure 5. Program-specific individual goal satisfaction ratings from Study 2.

suggested that there ought to be specific instructions within the online program to guide users to the book, such as indicating which pages were associated with a certain module and where the homework page was located. In general, participants did not utilize the “coffee break” chat rooms because of technical glitches, lack of interest, and scheduling conflicts.

Lastly, participants shared a sense of relief when learning about normal age-related memory changes experienced by most adults. They also tended to feel more in control of their memory, with one participant indicating that the best part of the program was “motivating me to take charge and that I shouldn’t be so quick to accept that losing some memory is unavoidable.” Additionally, participants consistently attributed their enjoyment and engagement to the many types of formats, games, activities, and homework.

Final iterations

Based on the feedback obtained throughout the agile development cycle, several final changes were made (e.g., adding to the FAQ document, dropping the “coffee break” chat rooms, making discussion board questions more open-ended). It was further decided that there would be increased moderation within the discussion boards, as having encouraging feedback, inviting participant responses, and redirecting to the goal of the topic at hand can increase engagement and provide an organized structure (Cudney & Weinert, 2000; Nahm et al., 2011). This can also be an opportunity for the facilitator to reinforce and elaborate on evidence-based information regarding memory and health, debunk any common misconceptions, and promote feelings of normalcy among participants (Pike et al., 2018).

Attrition in Study 2 was in part the result of technical obstacles, according to participant e-mails notifying the e-learning team. Despite addressing technical issues between Sub-studies 2a and 2b, the attrition rate remained at 50 per cent. The schedule was structured to release a module once a week, as long as the participant had completed the previous module. This resulted in a minimum requirement of 8 weeks of participation in the online program. Some participant feedback from open-ended questions indicated confusion surrounding when the module would be

released as well as concern from participants travelling without access to a computer. Further, participants may lose interest if there is a long wait period before the next module is released. Therefore, in future versions of the program, participants will be able to access the next module after completion of the preceding activities, which will allow participants to complete the program at their desired pace.

General Discussion

Development of Online Memory Programs

Over the coming years, there will undoubtedly be an increasing number of online health programs to serve the aging Canadian population. Although there are practice recommendations to tailor memory programs to the unique needs of older adults (Pike et al., 2018), this is the first article to provide a detailed description of a process that can be adapted to achieve such personalization. For example, Rebok, Tzuang, and Parisi (2020) did not provide information related to the development of their online memory training program, ACTIVE Memory Works™, and did not describe any piloting conducted prior to their randomized study. In a recent publication of a study protocol for the online memory program OPTIMiSe (Pike et al., 2021), there is mention of obtaining feedback from an “advisory committee” consisting of health experts and individuals from the target group. However, there is no indication of whether or how any feedback was integrated into the final protocol, and whether the program was pilot tested on any end users.

Overall, adapting an agile development cycle fostered collaboration and supported inclusion of important input from diverse stakeholders. We were open to the process of small changes and many iterations, and adopted a sense of flexibility with timelines and the number of piloting sessions needed to produce the final product. In addition, following the five clinical and translational phases offered a clear framework and structure to the project. Through this process, we provided evidence that our online memory program was designed to be user friendly and enjoyable, while demonstrating preliminary benefits associated with the

in-person version. In the future, researchers can consider including a user experience (UX) designer to facilitate the agile process and to video record piloting sessions in order to systematically code and evaluate users' experience, similarly to the methodology applied in Mansson et al.'s (2020) co-creation of a smart phone application with older adult end users. Although analytic data were monitored to track program progress in Study 2, future avenues for research include analyzing data to further investigate any areas of navigation difficulty and whether there are any associations between analytics (e.g., average length of time between the completion of various modules) and program benefits.

Following the successful completion of the T2 and T3 translational phases described in this article, future research is ongoing to conduct a wider implementation of the online memory program to evaluate true benefits within the community characteristic of the final T4 Translation to Population Health phase. Specifically, as a first step in this process, we are currently conducting a randomized controlled trial to evaluate benefits of participating in the program in comparison with a no-intervention control group (i.e., treatment as usual for healthy older adults; Petersen et al., 2018).

Limitations

Study 2 had 50 per cent attrition across the two sequential pilots. This was not unexpected, however, because online interventions tend to have higher rates of attrition than in-person interventions (Eysenbach, 2005; Kelders et al., 2012; Peels et al., 2012). High attrition rates may occur for a variety of reasons such as the fleeting or "surfing" culture of the Internet (Ahern, 2007). It may be that participants feel a greater sense of responsibility or investment when participating in person as there is more rapport established between intervention facilitators and other group members. According to Eysenbach (2005), attrition is "one of the fundamental characteristics and methodological challenges in the evaluation of eHealth applications" (p. 2). Understanding the reasons for participant attrition from online interventions and being able to predict or control such attrition is an emerging area of research. Although data were logged for participants who volunteered their reasons for leaving the study, we did not routinely inquire about the reasons for discontinuation in cases for which this was not offered. Therefore, this is an area in need of future systematic research; for example, using a feedback questionnaire designed to understand the reasons for discontinuing an online program. It would be interesting to explore what information can be used to understand group differences between individuals who drop out and those who complete an online intervention.

Another important limitation of the described agile development cycle is the extensive use of time and resources. From the first translational phase to the completion of Study 2 (T3 Translation to Patients phase), the project spanned more than 3 years and involved a large multidisciplinary team of researchers, clinicians, e-learning designers, patient advisors, research assistants, and administrative staff. The piloting described in Study 1 required the use of an on-site computer laboratory with headphones. These costs of course are an important consideration in undertaking such a project, but should be weighed against the potential monetary and non-monetary (e.g., poor user experience) costs of making post-implementation revisions if the project was not designed properly in the first place.

Conclusion

Considering the various limitations of in-person programs, there is growing interest in the development of online health interventions for older adults. Although there are clear practice recommendations to tailor online memory programs to the needs of older adults, our article is the first to describe the adoption of the agile development cycle with the clinical and translational phases framework to develop and pilot an online program while integrating feedback from older adults. Through this process, we were able to ensure that our online memory program was user friendly and enjoyable to use, and that it demonstrated targeted program benefits. Such a process can be adapted by others interested in offering online content to a target population.

Acknowledgments. We thank Derek Redmond for his assistance with the software development of the online program and all of the research assistants involved in assisting with data collection.

References

- AGE-WELL. (2020). Tech use by older Canadians for health, wellness and independence in the time of COVID-19. Retrieved 20 October 2020 from <https://agewell-nce.ca/wp-content/uploads/2020/09/OATechSurvey-SEP2020-FINAL.pdf>.
- Ahern, D. K. (2007). Challenges and opportunities of eHealth research. *American Journal of Preventive Medicine*, *32*(5), S75–S82. <https://doi.org/10.1016/j.amepre.2007.01.016>
- Astell, A., Dove, E., Morland, C., & Donovan, S. (2020). Using the TUNGSTEN approach to co-design DataDay: A self-management app for dementia. In R. Brankaert & G. Kenning (Eds.), *HCI and design in the context of dementia*, Human-Computer Interaction Series (pp. 171–185). Cham: Springer. https://doi.org/10.1007/978-3-030-32835-1_11
- Chang, J., McAllister, C., & McCaslin, R. (2015). Correlates of, and barriers to, internet use among older adults. *Journal of Gerontological Social Work*, *58*(1), 66–85. <https://doi.org/10.1080/01634372.2014.913754>
- Craik, F. I. M., & Rose, N. S. (2012). Memory encoding and aging: A neurocognitive perspective. *Neuroscience and Biobehavioral Reviews*, *36*(7), 1729–1739. <https://doi.org/10.1016/j.neubiorev.2011.11.007>
- Cudney, S. A., & Weinert, C. (2000). Computer-based support groups. Nursing in cyberspace. *Computers in Nursing*, *18*(1), 35–43.
- Davis, B. (2013). *Agile practices for waterfall projects: Shifting processes for competitive advantage*. Plantation, FL: J. Ross Publishing
- Doolittle, J. (2020). Jeremy Miller on waterfall versus agile. *IEEE Software*, *37*(4), 107–C3. <https://doi.org/10.1109/MS.2020.2987493>
- Eysenbach, G. (2005). The law of attrition. *Journal of Medical Internet Research*, *7*(1), e11. <https://doi.org/10.2196/jmir.7.1.e11>
- Fitzpatrick, A. L., Powe, N. R., Cooper, L. S., Ives, D. G., & Robbins, J. A. (2004). Barriers to health care access among the elderly and who perceives them. *American Journal of Public Health*, *94*(10), 1788–1794. <https://doi.org/10.2105/AJPH.94.10.1788>
- Flewelling, P. (2018). *The agile developer's handbook*. Packt Publishing.
- Harvard Catalyst. (2021). Pathfinder. [catalyst.harvard.edu](https://catalyst.harvard.edu/pathfinder/). Retrieved 9 May 2021 from <https://catalyst.harvard.edu/pathfinder/>.
- Hudes, R., Rich, J. B., Troyer, A. K., Yusupov, I., & Vandermorris, S. (2019). The impact of memory intervention programs on subjective outcomes in healthy older adults: A systematic review and meta-analysis. *Psychology and Aging*, *34*(4), 587–597. <https://doi.org/10.1037/pag0000340>
- Karbach, J., & Verhaeghen, P. (2014). Making working memory work: A meta-analysis of executive-control and working memory training in older adults. *Psychological Science*, *25*(11), 2027–2037. <https://doi.org/10.1177/0956797614548725>
- Kelders, S. M., Kok, R. N., Ossebaard, H. C., & van Gemert-Pijnen, J. E. W. C. (2012). Persuasive system design does matter: A systematic review of adherence to web-based interventions. *Journal of Medical Internet Research*, *14*(6), e152. <https://doi.org/10.2196/jmir.2104>

- Kueider, A. M., Parisi, J. M., Gross, A. L., & Rebok, G. W. (2012). Computerized cognitive training with older adults: A systematic review. *PLoS One*, *7*(7), e40588. <https://doi.org/10.1371/journal.pone.0040588>
- Ludden, G. D. S., van Rompay, T. J. L., Kelders, S. M., & van Gemert-Pijnen, J. E. W. C. (2015). How to increase reach and adherence of web-based interventions: A design research viewpoint. *Journal of Medical Internet Research*, *17*(7), e172. <https://doi.org/10.2196/jmir.4201>
- Mansson, L., Wiklund, M., Öhberg, F., Danielsson, K., & Sandlund, M. (2020). Co-creation with older adults to improve user-experience of a smartphone self-test application to assess balance function. *International Journal of Environmental Research and Public Health*, *17*(11), 3768. <https://doi.org/10.3390/ijerph17113768>
- Nahm, E.-S., Bausell, B., Resnick, B., Covington, B., Brennan, P. F., Mathews, R., & Park, J. H. (2011). Online research in older adults: Lessons learned from conducting an online randomized controlled trial. *Applied Nursing Research*, *24*(4), 269–275. <https://doi.org/10.1016/j.apnr.2009.09.004>
- Parikh, P. K., Troyer, A. K., Maione, A. M., & Murphy, K. J. (2016). The impact of memory change on daily life in normal aging and mild cognitive impairment. *The Gerontologist*, *56*(5), 877–885. <https://doi.org/10.1093/geront/gnv030>
- Peels, D. A., Bolman, C., Golsteijn, R. H., De, V. H., Mudde, A. N., van Stralen, M. M., & Lechner, L. (2012). Differences in reach and attrition between web-based and print-delivered tailored interventions among adults over 50 years of age: Clustered randomized trial. *Journal of Medical Internet Research*, *14*(6), e179. <https://doi.org/10.2196/jmir.2229>
- Petersen, R. C., Lopez, O., Armstrong, M. J., Getchius, T. S. D., Ganguli, M., Gloss, D., et al. (2018). Practice guideline update summary: Mild cognitive impairment: Report of the guideline development, dissemination, and implementation subcommittee of the American Academy of Neurology. *Neurology*, *90*(3), 126–135. <https://doi.org/10.1212/WNL.0000000000004826>
- Pike, K. E., Chong, M. S., Hume, C. H., Keech, B. J., Konjarski, M., Landolt, K. A., et al. (2018). Providing online memory interventions for older adults: A critical review and recommendations for development. *Australian Psychologist*, *53*(5), 367–376. <https://doi.org/10.1111/ap.12339>
- Pike, K. E., Moller, C. I., Bryant, C., Farrow, M., Dao, D. P., & Ellis, K. A. (2021). Online personalised training in memory strategies for everyday (OPTIMiSE) program for older adults with cognitive concerns: Pilot study protocol. *Journal of Alzheimer's Disease Reports*, *5*(1), 143–152. <https://doi.org/10.3233/ADR-200251>
- Rebok, G. W., Tzuang, M., & Parisi, J. M. (2020). Comparing web-based and classroom-based memory training for older adults: The ACTIVE memory works™ study. *The Journals of Gerontology. Series B, Psychological Sciences and Social Sciences*, *75*(6), 1132–1143. <https://doi.org/10.1093/geronb/gbz107>
- Rübsamen, N., Akmatov, M. K., Castell, S., Karch, A., & Mikolajczyk, R. T. (2017). Factors associated with attrition in a longitudinal online study: Results from the HaBIDS panel. *BMC Medical Research Methodology*, *17*(1), 1–11. <https://doi.org/10.1186/s12874-017-0408-3>
- Stoica, M., Ghilic-Micu, B., Mircea, M., & Uscatu, C. (2016). Analyzing agile development – from waterfall style to Scrumban. *Informatica Economica*, *20*(4), 5–14. <https://doi.org/10.12948/issn14531305/20.4.2016.01>
- Sung, N. S., Crowley, J., William F, Genel, M., Salber, P., Sandy, L., et al. (2003). Central challenges facing the national clinical research enterprise. *JAMA: The Journal of the American Medical Association*, *289*(10), 1278–1287. <https://doi.org/10.1001/jama.289.10.1278>
- Szilagyi, P. (2009). Translational Research and Pediatrics. *Academic Pediatrics*, *9*(2), 71–80. <https://doi.org/10.1016/j.acap.2008.11.002>
- Troyer, A. K. (2001). Improving memory knowledge, satisfaction, and functioning via an education and intervention program for older adults. *Aging, Neuropsychology, and Cognition*, *8*(4), 256–268. <https://doi.org/10.1076/anec.8.4.256.5642>
- Troyer, A. K., & Vandermorris, S. (2012). *Memory and aging program: Leader's manual*. Toronto: Baycrest Centre for Geriatric Care.
- Troyer, A. K., & Vandermorris, S. (2017). *Memory and aging program: Clinical guide*. Toronto: Baycrest Centre for Geriatric Care.
- Vandermorris, S., Au, A., Gardner, S., & Troyer, A. K. (2020). Initiation and maintenance of behaviour change to support memory and brain health in older adults: A randomized controlled trial. *Neuropsychological Rehabilitation*. <https://doi.org/10.1080/09602011.2020.1841656>
- Vandermorris, S., Davidson, S., Au, A., Sue, J., Fallah, S., & Troyer, A. (2017). “Accepting where I’m at” – A qualitative study of the mechanisms, benefits, and impact of a behavioral memory intervention for community-dwelling older adults. *Aging & Mental Health*, *21*(9), 895–901. <https://doi.org/10.1080/13607863.2016.1181709>
- Westfall, J. M., Mold, J., & Fagnan, L. (2007). Practice-based research—“Blue highways” on the NIH roadmap. *JAMA: The Journal of the American Medical Association*, *297*(4), 403–406. <https://doi.org/10.1001/jama.297.4.403>
- Wiegand, M. A., Troyer, A. K., Gojmerac, C., & Murphy, K. J. (2013). Facilitating change in health-related behaviors and intentions: A randomized controlled trial of a multidimensional memory program for older adults. *Aging & Mental Health*, *17*(7), 806–815. <https://doi.org/10.1080/13607863.2013.789000>